

REMARK ON CONSERVATION LAWS ASSOCIATED WITH NON-NOETHER SYMMETRIES

George Chavchanidze

Department of Theoretical Physics
A. Razmadze Institute Mathematics

1 Aleksidze Street, Ge 380093
Tbilisi, Georgia

e-mail:gch@rmi.acnet.ge

Abstract

In the present paper geometric aspects of relationship between non-Noether symmetries and conservation laws in Hamiltonian systems is discussed. It is shown that integrals of motion associated with continuous non-Noether symmetry are in involution whenever generator of the symmetry satisfies certain Yang-Baxter equation.

2000 Mathematical Subject Classification: 70H33, 70H06, 53Z05

In Hamiltonian systems time evolution of observables is governed by Hamilton's equation

$$\frac{d}{dt}f = \{h, f\} \quad (1)$$

here $\{, \}$ is Poisson bracket defined by some bivector field W satisfying condition

$$[W, W] = 0. \quad (2)$$

Namely

$$\{f, g\} = W(df \wedge dg) = L_{W(f)}g = -L_{W(g)}f \quad (3)$$

where $W(f)$ and $W(g)$ are Hamiltonian vector fields associated with f and g functions and L is Lie derivative. Skew symmetry of bivector field W provides skew symmetry of Poisson bracket, while (2) condition ensures Jacobi identity. We also assume that dynamical system is regular - bivector field has maximal rank (i. e. $W^n \neq 0$).

Each vector field E defines continuous one-parameter group of transformations $g_a = e^{aL_E}$ that acts on observables

$$g_a(f) = e^{aL_E}(f) = f + aL_E f + \frac{1}{2}a^2 L_E^2 f + \dots \quad (4)$$

Such a group of transformation is a symmetry of (1) Hamilton's equation whenever its generator E commutes with evolution operator $W(h) = \{h, \}$ i. e.

$$[E, W(h)] = 0. \quad (5)$$

Symmetry is said to be Noether (or Cartan) if its generator is Hamiltonian vector field (such a vector fields preserve W) and non-Noether (non-Cartan) whenever generator is non-Hamiltonian (in other words $[E, W] \neq 0$). Non-Noether symmetries give rise to a number of integrals of motion described in the following theorem [12] (see also [2] [3] [8] [10] [11])

Theorem: If the vector field E generates non-Noether symmetry, then the functions

$$Y^{(l)} = \frac{\hat{W}^l \wedge W^{n-l}}{W^n} \quad l = 1, 2, \dots, n \quad (6)$$

where $\hat{W} = [E, W]$ and n is half dimension of the phase space, are constant along solutions.

Proof: By definition

$$\hat{W}^l \wedge W^{n-l} = Y^{(l)} W^n \quad (7)$$

now let us take Lie derivative of this expression along the vector field $W(h)$.

$$[W(h), \hat{W}^l \wedge W^{n-l}] = \frac{d}{dt}(Y^{(l)})W^n + Y^{(l)}[W(h), W^n] \quad (8)$$

or

$$l[W(h), \hat{W}] \wedge \hat{W}^{l-1} \wedge W^{n-l} + (n-l)[W(h), W] \wedge \hat{W}^l \wedge W^{n-l-1} = \frac{d}{dt}(Y^{(l)})W^n + nY^{(l)}[W(h), W] \wedge W^{n-1} \quad (9)$$

but according to Liouville theorem Hamiltonian vector field preserves W i. e.

$$[W(h), W] = 0 \quad (10)$$

and as a result

$$[W(h), \hat{W}] = [W(h)[E, W]] = [W[W(h), E]] + [E[W, W(h)]] = 0 \quad (11)$$

since $[W(h), E] = 0$. So $\frac{d}{dt}Y^{(l)} = 0$.

Remark. Instead of conserved quantities $Y^{(1)}...Y^{(n)}$ solutions $c_1...c_n$ of the secular equation

$$(\hat{W} - cW)^n = 0 \quad (12)$$

could be associated with generator of symmetry. By expanding expression (12) it is easy to verify that $Y^{(l)}$ integrals of motion can be expressed by means of $c_1...c_n$ functions as follows

$$Y^{(l)} = \frac{(n-l)!l!}{n!} \sum_{i_p \neq i_s} c_{i_1} c_{i_2} ... c_{i_l} \quad (13)$$

So n integrals of motion are associated with each generator of non-Noether symmetry. According to Liouville-Arnold theorem Hamiltonian system is completely integrable if it possesses n functionally independent integrals of motion in involution (two functions f and g are said to be in involution if their Poisson bracket vanishes $\{f, g\} = 0$). Generally speaking conservation laws associated with symmetry might appear to be neither independent nor involutive. However they are in involution if generator of the symmetry satisfies Yang-Baxter type equation. Namely we have the following theorem.
Theorem. If the vector field E satisfies condition

$$[[E[E, W]]W] = 0 \quad (14)$$

then the functions (6) are in involution

$$\{Y^{(k)}, Y^{(l)}\} = 0 \quad (15)$$

Proof: First of all let us note that identity (2) satisfied by Poisson bivector field W is responsible for Liouville theorem

$$[W, W] = 0 \quad \leftrightarrow \quad L_{W(f)}W = [W(f), W] = 0 \quad (16)$$

By taking Lie derivative of the (2) we obtain another useful identity

$$L_E[W, W] = [E[W, W]] = [[E, W]W] + [W[E, W]] = 2[\hat{W}, W] = 0 \quad (17)$$

This identity gives rise to the following relation

$$[\hat{W}, W] = 0 \quad \leftrightarrow \quad [\hat{W}(f), W] = -[\hat{W}, W(f)] \quad (18)$$

an finally condition (14) ensures third identity

$$[\hat{W}, \hat{W}] = 0 \quad (19)$$

yielding Liouville theorem for $\hat{W}$

$$[\hat{W}, \hat{W}] = 0 \quad \leftrightarrow \quad [\hat{W}(f), \hat{W}] = 0 \quad (20)$$

Indeed

$$[\hat{W}, \hat{W}] = [[E, W]\hat{W}] = [[\hat{W}, E]W] = -[[E, \hat{W}]W] = -[[E[E, W]]W] = 0 \quad (21)$$

Now let us consider two different solution $c_i \neq c_j$ of (12) equation. By taking derivative of

$$(\hat{W} - c_i W)^n = 0 \quad (22)$$

equation along $W(c_j)$ and $\hat{W}(c_j)$ vector fields and using Liouville theorem for W and $\hat{W}$ bivectors we obtain the following relations

$$(\hat{W} - c_i W)^{n-1} (L_{W(c_j)} \hat{W} - \{c_j, c_i\} W) = 0 \quad (23)$$

and

$$(\hat{W} - c_i W)^{n-1} (c_i L_{\hat{W}(c_j)} W + \{c_j, c_i\}_\bullet W) = 0 \quad (24)$$

where

$$\{c_i, c_j\}_\bullet = \hat{W}(dc_i \wedge dc_j) \quad (25)$$

is Poisson bracket calculated by means of $\hat{W}$ bivector field. Now multiplying (23) by c_i subtracting (24) and using (18) identity gives rise to

$$(\{c_i, c_j\}_\bullet - c_j \{c_i, c_j\})(\hat{W} - c_i W)^{n-1} W = 0 \quad (26)$$

So either

$$\{c_i, c_j\}_\bullet - c_j \{c_i, c_j\} = 0 \quad (27)$$

or volume field $(\hat{W} - c_i W)^{n-1} W$ vanishes. In last case we can iterate (23-27) procedure yielding after n iterations $W^n = 0$ that according to our assumption (dynamical system is regular) is not true. As a result we have (27) and by simple exchange of indices $i \leftrightarrow j$ we get

$$\{c_i, c_j\}_\bullet - c_i \{c_i, c_j\} = 0 \quad (28)$$

Finally by comparing (27) and (28) we obtain that c_i functions are in involution with respect to both Poisson structures (since $c_i \neq c_j$)

$$\{c_i, c_j\}_\bullet = \{c_i, c_j\} = 0 \quad (29)$$

and according to (13) the same is true for $Y^{(l)}$ integrals of motion.

Note. Each generator of non-Noether symmetry satisfying equation (14) endows dynamical system with bi- Hamiltonian structure [1] [3] [4] [7] [9] - couple $(W, \hat{W})$ of compatible $([W, \hat{W}] = 0)$ Poisson $([W, W] = [\hat{W}, \hat{W}] = 0)$ bivector fields and give rise to bicomplex structure - [4][5] [6] [9] couple $(d = [W, \cdot], \tilde{d} = [\hat{W}, \cdot])$ of compatible $(\tilde{d}d + d\tilde{d} = 0)$ differential $(d^2 = \tilde{d}^2 = 0)$ operators on graded algebra of vector fields.

Example: To illustrate correspondence between non-Noether symmetries and conservation laws let us consider simple example (other examples can be found in [3]) of particle experiencing action of dissipating force. If friction depends linearly on velocity then time evolution in such a system is governed by Hamilton's equation (1) with

$$W = \sum_{i=1}^n p_i \partial_{p_i} \wedge \partial_{q_i} \quad (30)$$

and

$$h = \sum_{i=1}^n p_i + q_i \quad (31)$$

And evolution operator is

$$X = \sum_{i=1}^n p_i (\partial_{q_i} - \partial_{p_i}) \quad (32)$$

This simple Hamiltonian system possesses nontrivial symmetry - it is invariant under the following infinitesimal transformations

$$q_i \rightarrow q_i + \epsilon(p_i + q_i)^2 \quad (33)$$

generated by

$$E = \sum_{i=1}^n (p_i + q_i)^2 \partial_{q_i} \quad (34)$$

vector field. Clearly E satisfies commutation relation (5) and Yang-Baxter equation (14). Corresponding conserved quantities

$$c_i = p_i + q_i \quad (35)$$

are in involution.

1 Acknowledgements

Author is grateful to Z. Giunashvili for constructive discussions. This work was supported by INTAS (00-00561).

References

- [1] R. Brouzet, Sur quelques propriétés géométriques des variétés bihamiltoniennes, 1989 C. R. Acad. Sci. Paris 308, série I, 287-92
- [2] G. Chavchanidze, Non-Noether symmetries in singular dynamical systems, 2001 Georgian Math. J. 8 (2001) 027-032, math-ph/0105037
- [3] G. Chavchanidze, Bi-Hamiltonian structure as a shadow of non-Noether symmetry, 2001, math-ph/0106018 to appear in Georgian Math. J. (April 2003)
- [4] M. Crampin, W. Sarlet, G. Thompson, Bi-differential calculi and bi-Hamiltonian systems, 2000 J. of Phys. A: Math. Gen. 33 No. 22 L177-180

- [5] A. Dimakis, F. Müller-Hoissen, Bi-differential calculi and integrable models, 2000 J. of Phys. A: Math. Gen. 33 210-23
- [6] A. Dimakis, F. Müller-Hoissen, Bicomplexes and integrable models, 2000 nlin.SI/0006029
- [7] G. Falqui, F. Magri, M. Pedroni, The method of Poisson pairs in the theory of nonlinear PDEs, 2000 nlin.SI/0002009
- [8] F. González-Gascón, Geometric foundations of a new conservation law discovered by Hojman, 1994 J. of Phys. A: Math. Gen. 27 L59-60
- [9] P. Guha, A Note on Bidifferential Calculi and Bihamiltonian systems, 2000 IHÉS preprint M/64
- [10] S. Hojman, A new conservation law constructed without using either Lagrangians or Hamiltonians, 1992 J. of Phys. A: Math. Gen. 25 L291-295
- [11] M. Lutzky, Remarks on a recent theorem about conserved quantities, 1995 J. of Phys. A: Math. Gen. 28 L637-638
- [12] M. Lutzky, New derivation of a conserved quantity for Lagrangian systems, 1998 J. of Phys. A: Math. Gen. 15 L721-722